



THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Göran KARLSSON

Appl. No. 09/986,464

Filed: November 8, 2001

For: **Process for Preparing Latent
Antithrombin III**

Confirmation No. 5096

Art Unit: 1645

Examiner: To Be Assigned

Atty. Docket: 2081.0020001/EEF

**Claim For Priority Under 35 U.S.C. § 119(a)-(d) In Utility
Application**

Commissioner for Patents
Washington, D.C. 20231

Sir:

Priority under 35 U.S.C. § 119(a)-(d) is hereby claimed to the following priority document, filed in a foreign country within twelve (12) months prior to the filing of the above-referenced United States utility patent application:

Country	Priority Document Appl. No.	Filing Date
SWEDEN	0004086-5	11/08/2000

A certified copy of each listed priority document is submitted herewith. Prompt acknowledgment of this claim and submission is respectfully requested.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

Eldora Ellison Floyd
Attorney for Applicant
Registration No. 39,967

Date: 1/25/02

1100 New York Avenue, N.W.
Suite 600
Washington, D.C. 20005-3934
(202) 371-2600

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen



Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande Biovitrum AB, Stockholm SE
Applicant (s)

(21) Patentansökningsnummer 0004086-5
Patent application number

(86) Ingivningsdatum 2000-11-08
Date of filing

Stockholm, 2001-12-19

För Patent- och registreringsverket
For the Patent- and Registration Office


Hjordis Segerlund

Avgift
Fee 170:-

PREPARATION PROCESS**Field of the invention**

The present invention relates to a process for the preparation of latent antithrombin III.

5

Background of the invention

Antithrombin III (AT) is a plasma glycoprotein with a total molecular weight of 58.1 kDa (Lebing et al, Vox Sang. 67, 117-124, 1994), that inhibits serine proteases in the coagulation cascade and thus plays a major role in the regulation of blood clotting. Antithrombin III is an inhibitor of Factors IXa, Xa, XI and XIIa, as well as of thrombin. Thus, AT regulates clot formation in different stages of the coagulation cascade. A small decrease of AT content in the blood is associated with an increased risk of thromboembolism. Concentrates of AT are used in the prophylaxis and treatment of thromboembolic disorders in patients with acquired or hereditary antithrombin deficiency. In addition, it has been reported that AT has a function in many other processes of the human body, for example in angiogenesis and inflammatory responses. The function of AT in these physiological processes is not fully understood.

A particular form of antithrombin III, which was first characterised by Wardell et al (Biochemistry 36, 13133-13142, 1997), is known as the latent form (L-AT). L-AT and a selectively elastase cleaved variant have been shown to possess a strong antiangiogenic activity, and also to suppress tumor growth in mice that have been injected subcutaneously with a human neuroblastoma cell line (O'Reilly et al, Science 285, 1926-1928, 1999, and WO 00/20026). Hence, L-AT must be considered a potential human anti-cancer drug. However, clinical evaluation of this potential drug remains to be performed.

Purification of AT with affinity chromatography is done using purified heparin as solid phase bound ligand, as is known in the art. Miller-Andersson et al (Thrombosis Research 5, 439-452, 1974) discloses the use of heparin-Sepharose to purify human AT. This chromatographical system has also been useful for the separation between AT and L-AT, where the decreased affinity of heparin for L-AT relative to AT makes it possible to resolve the two components, as described by Chang and Harper (Thrombosis and Haemostasis 77, 323-328, 1997).

Induction of the latent form of AT has previously been performed as described by Wardell et al (*supra*), who obtained 50-60% L-AT by incubating AT in 0.25 M citrate, 10 mM Tris/HCl, pH 7.4, for 15 h in 60°C.

Upon incubation of native antithrombin III at 60°C in medium or buffer only, aggregates of polymerised protein are often formed. The presence of these aggregates is detrimental to a high yield of latent antithrombin III, and should be avoided as far as possible.

An object of the present invention is then to provide a process for the preparation of latent antithrombin III, L-AT, from a solution of native antithrombin III, AT, which process gives a higher yield of the desired product than the prior art process.

A further object of the invention is to provide a process for the preparation of L-AT from AT, wherein the production of aggregates of AT polymers is kept to a minimum.

Another object of the invention is to provide such a process for the conversion of AT to L-AT, in which commonly available reagents and buffer solutions are used, and which is performed in vitro.

Still another object of the invention is to provide a process for the preparation of L-AT from AT, which is readily scaled up for industrial production of L-AT.

Summary of the invention

The aforementioned and other objects of the invention are met by a process as defined in the claims. Thus, a process is provided, which comprises incubation of a solution of native antithrombin III in the presence of sulphate ions and a buffer selected from Good's zwitterionic buffers. It has surprisingly been found that these incubation conditions makes possible the recovery of L-AT from the process in yields that are substantially higher than those obtained by methods of the prior art (notably the citrate conditions of Wardell et al), while avoiding possible aggregation problems.

Figure legends

Figure 1: Heparin affinity chromatography of anti-thrombin using a sodium chloride gradient, 0-2 M (5-60 min). The injected amount of protein was 100 µg for sample A-B, and 150 µg for sample C-D. All samples were incubated at 60°C for 16 h, except for the reference AT sample A, which was not heat-treated (sample 7 in the example). Sample B (sample 6 in the example) was incubated according to Wardell, ie in 0.5 M citrate. Samples C (sample 2 in the example) and D (sample 1 in the example) were incubated in 5 mM HEPES, pH 7.4, with 0.9 and 0.8 M ammonium sulphate respectively. Integration of the low affinity heparin-binding peak, eluting at 22 min, gave 44%, 71% and 89% of the total integrated area for samples B, C, and D, respectively. Native AT eluted at 39 min.

Figure 2: Native electrophoresis of antithrombin samples, using 12.5% polyacrylamide in a homogeneous gel. The amount of sample was 0.5 µg protein/lane, and the gels were silver-stained after running. All samples, except for lane 7, were incubated in 60°C for 16 h.

Lane 1) 5 mM HEPES, 0.8 M ammonium sulphate, pH 7.4
 Lane 2) 5 mM HEPES, 0.9 M ammonium sulphate, pH 7.4
 Lane 3) 5 mM HEPES, 1.1 M ammonium sulphate, pH 7.4
 Lane 4) 5 mM HEPES, 1.4 M ammonium sulphate, pH 7.4

- Lane 5) 5 mM HEPES, 2.0 M ammonium sulphate, pH 7.4
Lane 6) 10 mmol Tris/HCl, 0.5 M trisodium citrate, pH 7.4 (according to Wardell et al. 1997)
Lane 7) Reference AT sample, not heat-treated
5 Lane 8) 25 mM sodium phosphate, 100 mM sodium chloride, pH 7.4
Lane 9) 25 mM HEPES, 0.8 M ammonium sulphate, pH 7.4
Lane 10) 5 mM HEPES, 0.5 M ammonium sulphate, pH 7.4
Lane 11) 5 mM HEPES, 2.0 M ammonium sulphate, pH 7.4
10 Lane 12) 5 mM HEPES, 0.8 M ammonium sulphate, pH 7.0
All lane numbers correspond to the sample numbers listed in the example below.

Detailed description of the invention

- 15 The invention provides a process for the preparation of latent antithrombin III (referred to as L-AT), starting from a solution of antithrombin III in its native form (referred to as AT). AT can be isolated from blood plasma by heparin-Sepharose chromatography as has been
20 described. According to the invention, the AT is then incubated in the presence of sulphate ions and a buffer. The incubation temperature and duration can be readily determined by the skilled person, but normal pasteurisation conditions, such as a temperature of about 60°C during about 16 hours, have been found to work well.

- 25 The sulphate ions are preferably provided in the form of a sulphate salt. Here, the use of an alkali metal sulphate, an alkaline earth sulphate or ammonium sulphate is preferred. Especially preferred is the use of ammonium
30 sulphate. A suitable concentration of sulphate ions in the process according to the invention lies in the range from 0.5 to 2.0 M, preferably from 0.7 to 1 M, a concentration between 0.8 and 0.9 M being most preferred.

- 35 Another component of the incubation mixture is a buffer selected from Good's zwitterionic buffers (Good et al, Biochemistry 5, 467-477, 1966). Which of the indicated buffers to use in the process of the invention can

be determined without undue experimentation, keeping in mind that the buffer should fulfil most or all of the following requirements: it should exhibit a pK_a value of between about 6 and about 9, a maximum solubility in water and a minimum solubility in other solvents, produce a minimum of salt effects, be stable at the experimental conditions used, and not absorb light in the visible or ultraviolet spectral regions (so as not to interfere with spectrophotometric measurements). Good's zwitterionic buffers, including buffers such as HEPES, MES and PIPES, typically present the desired characteristics. The use of HEPES is particularly preferred in the process according to the invention. The widely used Tris buffer is unsuitable for the purposes of the invention. Preferred buffer concentrations are somewhat dependent on the buffer chosen, but typically lie in the range from 1 to 25 mM, more preferably from 2.5 to 10 mM, most preferably from 4 to 6 mM.

As indicated above, the pH of the incubation reaction should lie between pH 6 and pH 9, preferably between pH 7 and pH 8, most preferably between pH 7.4 and pH 7.6.

Following the incubation of AT under the conditions outlined above, separation of the L-AT thus obtained from remaining AT is preferably performed using heparin affinity chromatography. The L-AT exhibits a lower binding affinity to heparin than AT, eluting substantially faster and enabling easy separation of the two forms of anti-thrombin III.

The preparation of L-AT thus obtained is advantageously subjected to treatment for the inactivation or removal of pathogens, particularly in the form of viruses and prions. This can be done in any stage of the process using one of several methods for inactivation or removal known in the art, or combinations of such methods. Examples of such methods include chemical inactivation, heat inactivation, light inactivation, microwave inactivation and nano-filtration removal. A dead-end filtration proce-

dure with a high salt content, like that described in WO96/00237, is particularly preferred, alone or in combination with other procedures. The removal and inactivation of pathogens can also be performed when the antithrombin III molecules are in the native state, before conversion to L-AT.

The invention is further illustrated by the following, non-limiting example.

10 **EXAMPLE**

A laboratory sample of AT, > 95% pure, was obtained from Plasma Products, Pharmacia, Stockholm, Sweden. This sample was prepared according to known methods (Miller-Andersson et al, *supra*) and used for induction of the latent form of antithrombin.

Preparation of L-AT

The laboratory sample of AT was transferred to the following solutions:

- 20 Sample 1) 5 mM HEPES, 0.8 M ammonium sulphate, pH 7.4
- Sample 2) 5 mM HEPES, 0.9 M ammonium sulphate, pH 7.4
- Sample 3) 5 mM HEPES, 1.1 M ammonium sulphate, pH 7.4
- Sample 4) 5 mM HEPES, 1.4 M ammonium sulphate, pH 7.4
- 25 Sample 5) 5 mM HEPES, 2.0 M ammonium sulphate, pH 7.4
- Sample 6) 10 mmol Tris/HCl, 0.5 M trisodium citrate, pH 7.4 (according to Wardell et al. 1997)
- Sample 7-8) 25 mM sodium phosphate, 100 mM sodium chloride, pH 7.4
- 30 Sample 9) 25 mM HEPES, 0.8 M ammonium sulphate, pH 7.4
- Sample 10) 5 mM HEPES, 0.5 M ammonium sulphate, pH 7.4
- Sample 11) 5 mM HEPES, 2.0 M ammonium sulphate, pH 7.4
- Sample 12) 5 mM HEPES, 0.8 M ammonium sulphate, pH 7.0
- Sample 13) 5 mM HEPES, 0.8 M ammonium sulphate, pH 7.8

35

All buffers listed above were adjusted to the desired pH at room temperature; 1 M HCl was used for ad-

justment of sample 6, while 1 M sodium hydroxide was used for pH adjustment of all other samples.

AT at a final concentration of 6 mg/ml was incubated in the solutions (samples 1-13) in glass tubes for 16 h at 60°C (except for sample 7, which was kept in a fridge at about 8°C) and transferred to a solution containing 50 mM Tris/HCl, 50 mM sodium chloride, pH 7.4, using small gelfiltration columns (NAP-5 Amersham Pharmacia Biotech, Uppsala, Sweden).

The formation of L-AT in the samples was analysed by heparin affinity chromatography, and the presence of aggregates was analysed by native electrophoresis.

Heparin affinity chromatography

This method was performed based on Chang and Harper (*supra*). A HPLC equipped with an TSK Heparin® column (Toshaas, Stuttgart, Germany, 7.5 i.d. x 75 mm, 10 µm, 1000 Å) was used. Eluting buffers were 20 mM Tris/HCl buffer, pH 7.4 (buffer A) and 2 M sodium chloride in 20 mM Tris/HCl buffer, pH 7.4 (buffer B). A linear gradient was run (0-5 min of 0% B, 5-60 min 0-100% B, 60-90 min 0% B). The flow rate was 0.4 ml/min and detection was carried out by measuring the absorbance at 280 nm.

Native polyacrylamide gel electrophoresis

Electrophoresis was performed using a 12.5% polyacrylamide homogeneous Phast® gel (Amersham Pharmacia Biotech, Uppsala, Sweden) employing the recommended running parameters. 0.5 µg protein in 1 µl was loaded in each lane. A diamino silver staining was performed according to the booklet from Pharmacia & Upjohn (Phast System™, Technical Note No 2, Two-dimensional electrophoresis with PhastGel™ separation media, Pharmacia LKB Biotechnology AB, Uppsala, Sweden), except that use was made of a slightly stronger fixation solution, containing 50% ethanol, 10% acetic acid and 40% water.

Antithrombin activity

Sample 2 (incubation in 0.9 M ammonium sulphate) was analysed regarding biological AT activity with the thrombin chromogenic peptide substrate (S-2238) (Chromogenix, Molndal, Sweden), according to Handeland et al. (Scand J. Haematol. 31, 427-436, 1983). The assay solution consisted of thrombin, heparin, chromogenic substrate and sample, and the response after incubation was recorded as a change in absorbance at 405 nm.

Results

Heparin affinity chromatography gave elution of native AT at 39 min (about 0.9 M sodium chloride) and the main latent peak eluted at 22 min (about 0.3 M sodium chloride) (figure 1A-1B). Integration of the low heparin-binding peak indicated a yield of 44% (figure 1B) for the sample prepared according to Wardell's method (sample 6), while incubation in 0.9 and 0.8 M ammonium sulphate (samples 2 and 1, respectively) yielded 71% and 89% respectively of the total integrated area (figure 1C-1D). Table 1 shows that the percentage of formed L-AT decreases at increased concentration of ammonium sulphate/HEPES or at a higher pH value.

Native electrophoresis of AT incubated at 60°C in phosphate/NaCl (sample 8) gave a strong formation of aggregates, and only a minor part of the protein remained in the monomeric form (figure 2, lane 8). AT incubated according to Wardell (figure 2, lane 6), as well as the not incubated AT (figure 2, lane 7), gave no aggregates. Incubation in 0.5 M ammonium sulphate (sample 10) induced a strong aggregation (figure 2, lane 10), while 0.8 M (sample 1) only gave a minor part of aggregates (figure 2, lane 1). Ammonium sulphate at a concentration of 0.9 - 2.0 M (samples 2-5) resulted in no visible aggregates (figure 2, lanes 2-5). At pH 7.0, a lot of aggregates were observed (figure 2, lane 12), while a pH of 7.8 gave a smaller amount of aggregates (data not shown).

Antithrombin activity assay on sample 2 (with 0.9 M ammonium sulphate) showed that 34% of the original specific activity remained; this should be compared with the 29% yield of high affinity heparin-binding AT upon analysis of the same sample by affinity chromatography (Table 1).

Table 1: Heparin affinity chromatography. Formation of L-AT in various sample buffers after 16 h incubation in 60 °C.

Incubation solution	Sample no ¹	% AT with low heparin affinity
10 mmol Tris/HCl, 0.5 M citrate, pH 7.4 (Wardell)	6	44*
5 mM Hepes, 0.5 M ammonium sulfate, pH 7.4	10	99
5 mM Hepes, 0.8 M ammonium sulfate, pH 7.4	1	89
5 mM Hepes, 0.9 M ammonium sulfate, pH 7.4	2	71*
5 mM Hepes, 1.1 M ammonium sulfate, pH 7.4	3	56*
5 mM Hepes, 1.4 M ammonium sulfate, pH 7.4	4	49*
5 mM Hepes, 2.0 M ammonium sulfate, pH 7.4	5	48*
25 mM Hepes, 0.8 M ammonium sulfate, pH 7.4	9	70
5 mM Hepes, 0.8 M ammonium sulfate, pH 7.0	12	99
5 mM Hepes, 0.8 M ammonium sulfate, pH 7.8	13	65

¹ According to the example

* No visible aggregates when analysed by native electrophoresis (see Fig. 2)

15 Experimental conclusions

By incubation of AT in 5 mM HEPES, pH 7.4, containing 0.8 or 0.9 M ammonium sulphate in 60°C for 16 h, about 85-90% and 70-75% respectively of AT was transformed to the latent form. Native electrophoresis showed a small part of aggregates at 0.8 M ammonium sulphate and no visible aggregates at 0.9 M. In a purification procedure, such small amounts of aggregate can be easily removed by gel filtration or similar techniques.

The optimal concentration for the conversion of AT to L-AT using ammonium sulphate is 0.8-0.9 M. The conversion will also yield good results between 0.7 and 1 M, and some results between 0.5 and 2.0 M. For formation of L-AT, a process using 0.5-2.0 M ammonium sulphate, preferably 0.8-0.9 M, and up to 25 mM HEPES, preferably not more than 10 mM, at a pH near 7.4 has been found to give the most pleasing results. The percentage of L-AT formed will decrease at a higher concentration of ammonium sulphate/HEPES or at a higher pH value. In addition, for the prevention of formation of aggregates, it is necessary not to use too low an ammonium sulphate concentration or too low a pH value.

1108

CLAIMS

1. A process for the preparation of latent anti-thrombin III, comprising incubation of a solution of native antithrombin III in the presence of sulphate ions and a buffer selected from Good's zwitterionic buffers.
2. A process according to claim 1, wherein said sulphate ions are provided as a salt selected from ammonium sulphate, alkali metal sulphates and alkaline earth sulphates.
3. A process according to claim 2, wherein said sulphate salt is ammonium sulphate.
4. A process according to any one of claims 1-3, wherein the concentration of said sulphate ions is within the range from 0.5 to 2.0 M.
5. A process according to claim 4, wherein the sulphate ion concentration is within the range from 0.7 to 1 M.
6. A process according to claim 5, wherein the sulphate ion concentration is within the range from 0.8 to 0.9 M.
7. A process according to any one of claims 1-6, wherein said buffer comprises a HEPES buffer.
8. A process according to any one of claims 1-7, wherein the concentration of said buffer is within the range from 1 to 25 mM.
9. A process according to claim 8, wherein the buffer concentration is within the range from 2.5 to 10 mM.
10. A process according to claim 9, wherein the buffer concentration is within the range from 4 to 6 mM.
11. A process according to any one of claims 1-10, wherein the pH value is within the range from 6 to 9.
12. A process according to claim 11, wherein the pH value is within the range from 7 to 8.
13. A process according to claim 12, wherein the pH value is within the range from 7.4 to 7.6.

14. A process according to any one of claims 1-13, further including a treatment for inactivation or removal of pathogens, especially viruses and prions.

5 15. A process according to claim 1, wherein the native antithrombin III has been treated for inactivation or removal of pathogens, especially viruses and prions.

10 16. A process according to claim 14 or 15, wherein said treatment comprises one of, or a combination of, methods selected from chemical inactivation, heat inactivation, light inactivation, microwave inactivation and nano-filtration removal.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
219

ABSTRACT

A process for the preparation of latent antithrombin III is provided. The process comprises incubation of a solution of native antithrombin III in the presence of sulphate ions and a buffer selected from Good's zwitterionic buffers.

1/1

Absorbance (280 nm)




Fig. 1

33070000

2/2



Fig. 2